

# PROCEEDINGS OF SPIE

[SPIDigitalLibrary.org/conference-proceedings-of-spie](https://SPIDigitalLibrary.org/conference-proceedings-of-spie)

## A review of swarmalators and their potential in bio-inspired computing

Kevin O'Keeffe, Christian Bettstetter

Kevin O'Keeffe, Christian Bettstetter, "A review of swarmalators and their potential in bio-inspired computing," Proc. SPIE 10982, Micro- and Nanotechnology Sensors, Systems, and Applications XI, 109822E (13 May 2019); doi: 10.1117/12.2518682

**SPIE.**

Event: SPIE Defense + Commercial Sensing, 2019, Baltimore, Maryland, United States

# A review of swarmalators and their potential in bio-inspired computing

Kevin O’Keeffe<sup>a</sup> and Christian Bettstetter<sup>b</sup>

<sup>a</sup>Senseable City Lab, Massachusetts Institute of Technology, Cambridge, MA 02139, United States.

<sup>b</sup>Institute of Networked and Embedded Systems, University of Klagenfurt, 9020, Austria

## ABSTRACT

From fireflies to heart cells, many systems in Nature show the remarkable ability to spontaneously fall into synchrony. By imitating Nature’s success at self-synchronizing, scientists have designed cost-effective methods to achieve synchrony in the lab, with applications ranging from wireless sensor networks to radio transmission. A similar story has occurred in the study of swarms, where inspiration from the behavior flocks of birds and schools of fish has led to low-footprint algorithms for multi-robot systems. Here, we continue this ‘bio-inspired’ tradition, by speculating on the technological benefit of fusing swarming with synchronization. The subject of recent theoretical work, minimal models of so-called ‘swarmalator’ systems exhibit rich spatiotemporal patterns, hinting at utility in ‘bottom-up’ robotic swarms. We review the theoretical work on swarmalators, identify possible realizations in Nature, and discuss their potential applications in technology.

**Keywords:** Synchronization, swarming, bio-inspired computing.

## 1. INTRODUCTION

In 1967 Winfree proposed a model for coupled oscillators that spontaneously synchronize.<sup>1</sup> Beyond a critical coupling strength, the oscillators overcome the disordering effect of the dissimilarities in their natural frequencies and spontaneously lock their cycles. Kuramoto later simplified Winfree’s model and solved it exactly.<sup>2</sup> Since then, the study of sync has matured into a vibrant field.<sup>3–5</sup> On the theoretical side, theorists have twisted Kuramoto’s model in various ways resulting in rich phenomena like glassy behavior<sup>6–9</sup> and chimeras states.<sup>10–12</sup> On the applied side, coupled oscillators have found use in neurobiology,<sup>13–16</sup> cardiac dynamics,<sup>17–19</sup> and the bunching of school buses.<sup>20</sup> An interesting application of sync is the design of ‘bio-inspired’ algorithms. Here, by aping the simplicity of coupled oscillator models, researchers have devised resource-efficient algorithms to procure synchrony in the lab, useful for networked computing and robotics.<sup>21–24</sup>

A story with many parallels to the sync story has evolved in the study of swarms. In 1995 Vicsek proposed a simple model of swarming agents,<sup>25</sup> which – like the sync transition of coupled oscillators – showed a transition from disorder to order: beyond a critical coupling strength, the agents switched from a gas-like, incoherent state to one in which the agents moved as a coherent flock. The novelty of the out-of-equilibrium nature of the flocking transition – the system is out of equilibrium since agents constantly consume energy to propel themselves – piqued the minds of physicists and other theorists, which in turn helped give rise to the field of active matter,<sup>26–28</sup> a field perhaps more vibrant than the field of synchronization. In a final mirroring of the sync story, the bio-inspired community has also mimicked the minimalism of Vicsek’s and other models of swarming to design novel algorithms for optimization<sup>29–34</sup> and robotic swarms.<sup>33,35–38</sup>

These stories demonstrate swarming and synchronization are intimately related. In a sense, the two effects are ‘spatiotemporal opposites’: in synchronization the units self-organize in time but not in space; in swarming the units self-organize in space but not in time. Given this conceptual twinship, it is natural to wonder about the possibility of units which can self-organize in both space and time – that is, to wonder how swarming and synchronization might interact. And more pertinent to this work, to wonder how a mix of swarming and synchronization could be useful in technology.

---

Further author information: (Send correspondence to Kevin O’Keeffe; E-mail: [kokeeffe@mit.edu](mailto:kokeeffe@mit.edu))

Micro- and Nanotechnology Sensors, Systems, and Applications XI, edited by Thomas George,  
M. Saif Islam, Proc. of SPIE Vol. 10982, 109822E · © 2019 SPIE  
CCC code: 0277-786X/19/\$18 · doi: 10.1117/12.2518682

Several researchers have started to address these questions by analyzing systems that both sync and swarm. Von Brecht and Uminsky have generalized swarming particles by endowing them with an internal polarization vector,<sup>39</sup> equivalent to an oscillator's phase. Others have attacked the problem from the other direction, by considering synchronizing oscillators able to move around in space.<sup>40-43</sup> In these studies, however, the oscillators' movements affect their phase dynamics, but not the other way around; thus, the interaction between swarming and synchronization is only one-way. Two-way interaction between swarming and synchronization has also been considered: The pioneering work is the Iwasa-Tanaka model of chemotactic oscillators,<sup>44,45</sup> i.e., oscillators which interact through a background diffusing chemical. More recent works were carried out by Starnini et al,<sup>46</sup> Belovs et al,<sup>47</sup> and O'Keeffe et al<sup>48</sup> who proposed 'bottom-up' toy models without reference to a background medium. O'Keeffe et al called the elements of their systems 'swarmalators', to capture their twin identities as swarming oscillators, and to distinguish them from the mobile oscillators mentioned above for which the coupling between swarming and synchronization is unidirectional.

This paper reviews research on swarmalators and other systems that mix swarming and synchronization. Our main motivation is to identify if the interplay of sync and swarming can be useful for bio-inspired computing and related engineering domains. We outline the theoretical work on swarmalators, experimental realizations, and finally conjecture on their technological utility.

## 2. MODELS COMBINING SWARMING AND SYNCHRONIZATION

To combine synchronization with swarming, we first define what we mean by swarming. While to our knowledge there is no universally agreed on definition, swarming systems typically have at least one of two key features: (i) aggregation, arising from a balance between the units' mutual attraction and repulsion and (ii) alignment, referring to the units' tendency to align their orientation in space and move in a flock. Synchronization can thus be combined with either aggregation alone, alignment alone, or with both aggregation and alignment. In what follows, we present classes of model based on these three ways to combine sync and swarming.

### 2.1 Aggregation and Synchronization

The paradigmatic model of biological aggregation has the form

$$\dot{\mathbf{x}}_i = \frac{1}{N} \sum_{j \neq i}^N \mathbf{I}_{\text{att}}(\mathbf{x}_j - \mathbf{x}_i) - \mathbf{I}_{\text{rep}}(\mathbf{x}_j - \mathbf{x}_i), \quad (1)$$

where  $\mathbf{x}_i \in \mathbb{R}^d$  (usually with  $d \leq 3$ ) is the  $i$ -th particle's position,  $\mathbf{I}_{\text{att}}$  captures the attraction between particles, and  $\mathbf{I}_{\text{rep}}$  captures the repulsion between them. The competition between  $\mathbf{I}_{\text{att}}$  and  $\mathbf{I}_{\text{rep}}$  gives rise to congregations of particles with sharp boundaries, in accordance with many biological systems (see<sup>49,50</sup> for a review).

The paradigmatic model in synchronization is the Kuramoto model.<sup>51</sup>

$$\dot{\theta}_i = \omega_i + \frac{K}{N} \sum_{j \neq i}^N \sin(\theta_j - \theta_i). \quad (2)$$

Here  $\theta_i \in S^1$  and  $\omega_i$  are the phase and natural frequency of the  $i$ -th oscillator. The sine term captures the oscillators' coupling, where  $K > 0$  means oscillators tend to synchronize, and  $K < 0$  means oscillators tend to desynchronize. As mentioned, beyond a critical coupling strength  $K_c$ , a fraction of oscillators overcome the disordering effects imposed by their distributed natural frequencies  $\omega_i$  and spontaneously synchronize.

To study the co-action of synchronization and aggregation, a natural strategy would be to stitch the aggregation model (1) and the Kuramoto model (2) together. This was the approach taken by O'Keeffe et al<sup>48</sup> who proposed the following swarmalator model:

$$\dot{\mathbf{x}}_i = \frac{1}{N} \sum_{j \neq i}^N \left[ \frac{\mathbf{x}_j - \mathbf{x}_i}{|\mathbf{x}_j - \mathbf{x}_i|} \left( 1 + J \cos(\theta_j - \theta_i) \right) - \frac{\mathbf{x}_j - \mathbf{x}_i}{|\mathbf{x}_j - \mathbf{x}_i|^2} \right] \quad (3)$$

$$\dot{\theta}_i = \omega_i + \frac{K}{N} \sum_{j \neq i}^N \frac{\sin(\theta_j - \theta_i)}{|\mathbf{x}_j - \mathbf{x}_i|}. \quad (4)$$




Figure 1. **Swarmalator states.** Scatter plots in the  $(x, y)$  plane, where the swarmalators are colored according to their phase. (a) Static sync for  $(J, K) = (0.1, 1)$ . (b) Static async  $(J, K) = (0.1, -1)$ . (c) Static phase wave  $(J, K) = (1, 0)$ . (d) Splintered phase wave  $(J, K) = (1, -0.1)$ . (e) Active phase wave  $(J, K) = (1, -0.75)$ .

Equation (3) models phase-dependent aggregation and Equation (4) models position-dependent synchronization. The interaction between the space and phase dynamics is captured by the term  $1 + J \cos(\theta_j - \theta_i)$ . If  $J > 0$ , “like attracts like”: swarmalators are preferentially attracted to other swarmalators with the same phase, while  $J < 0$  indicates the opposite. For simplicity the authors considered identical swarmalators  $\omega_i = \omega$ , and by a change of reference they set  $\omega = 0$ .

The swarmalator model exhibits five long-term collective states. Figure 1 showcases these states as scatter plots in the  $(x, y)$  plane, where swarmalators are represented by dots and the color of each dot represents the swarmalator’s phase  $\theta$  (color, recall, can be mapped to  $S^1$  and so can be used to represent swarmalators’ phases). The parameter dependence of these states are encapsulated in the phase diagram shown in Figure 1. The first three states, named the *static sync*, *static async*, and *static phase wave*, are – as their names suggest – static in the sense that the individual swarmalators are stationary in both position and phase. This stationarity allows the density of these swarmalators  $\rho(\mathbf{x}, \theta)$  in these states to be constructed explicitly (the density  $\rho(\mathbf{x}, \theta)$  is interpreted in the Eulerian sense, so that  $\rho(\mathbf{x} + d\mathbf{x}, \theta + d\theta)$  gives the fraction of swarmalators with positions between  $\mathbf{x}$  and  $\mathbf{x} + d\mathbf{x}$  and phases between  $\theta$  and  $\theta + d\theta$ ). In the remaining *splintered phase wave* and *active phase waves* states swarmalators are no longer stationary. In the splintered phase wave state swarmalators execute small oscillation in both space and phase within each cluster. In the active phase wave the swarmalators split into counter-rotating groups – in both space and phase – so that  $\langle \dot{\theta}_i \rangle = \langle \dot{\phi}_i \rangle = 0$ , where  $\phi_i = \arctan(y_i/x_i)$  is the spatial angle of the  $i$ -th oscillator and angle brackets denote population averages. The conservation of these two quantities follows from the governing equations; the pairwise terms are odd and thus cancel under summation. Three-dimensional analogues of the five collective states were also reported.

The stability properties of the static async state are unusual. Via a linear stability analysis in density space, an integral equation for the eigenvalues  $\lambda$  was derived. Yet numerical solutions to this integral equation, in the parameter regime where the static async state should be stable, produced a leading eigenvalue so small in magnitude that its sign could not be determined reliably. Thus, the stability of the state could not be analytically confirmed. There is however a parameter value at which the magnitude of  $\lambda$  increases sharply, which was used to derive a pseudo critical parameter value  $K_c \approx -1.2J$  marking the apparent (i.e. as observed in simulations) destabilization of the state. Nevertheless, the true stability of the static async state remains a puzzle.

Some extensions of the work in<sup>48</sup> have been carried out. One addresses the conspicuously empty upper right quadrant in the  $(J, K)$  parameter plane in Figure 2, where only the trivial static sync state appears. By adding phase noise to (4), Hong discovered<sup>52</sup> the active phase wave exists for  $K > 0$ . The splintered phase wave however was not observed. O’Keeffe, Evers, and Kolokolnikov<sup>53</sup> have extended (3) by allowing phase similarity to affect the spatial repulsion term, as well as the spatial attraction term (i.e. they multiply the second term in (3) by a term  $1 + J_2 \cos(\theta_j - \theta_i)$ ). This led to the emergence of ring states, an example of which is depicted in Figure 3. They constructed and analyzed the stability of these ring states explicitly for a population of given size  $N$ . Analytic results for arbitrary  $N$  are potentially useful for robotic swarms, which presumably are realized in the small  $N$  limit. Another offshoot of this analysis was a heuristic method to predict the number of clusters which form in the splintered phase wave state; they viewed each cluster of synchronized swarmalators as one giant swarmalator which let them re-imagine the splintered phase wave state as a ring, allowing them to leverage their analysis. A precise description of the number of clusters formed is an open problem.




Figure 2. **Phase diagram.** Locations of states of the model defined by equations (3) and (4) in the  $(J, K)$  plane. The straight line separating the static async and active phase wave states is a semi-analytic approximation given by  $J \approx 1.2K$  – see Eq. (18) in.<sup>48</sup> Black and red dots show simulation data. The red dashed line simply connects the red dots and was included to make the boundary visually clearer. Note, this figure has been reproduced from.<sup>48</sup>




Figure 3. **Ring phase state.** Swarmalators are represented by colored dots in the  $(x, y)$  plane where the color of each dot represents its phase. The state is found by numerically integrating the governing equations in<sup>53</sup> with  $(J_1, J_2, K, N) = (0, 0.8, 0, 100)$ .

**Iwasa-Tanaka model.** Iwasa and Tanaka proposed and studied a ‘swarm-oscillator’ model in a series of papers.<sup>44, 54–57</sup> The inspiration for their work comes from chemotactic oscillators, i.e., oscillators moving around in a diffusing chemical which mediates their interactions. They began with the general model

$$\dot{\mathbf{X}}_i(t) = f(\mathbf{X}_i) + kg(S(\mathbf{r}_i, t)) \quad (5)$$

$$m\ddot{\mathbf{r}}_i(t) = -\gamma\dot{\mathbf{r}}_i - \sigma(\mathbf{X}_i)S \quad (6)$$

$$\tau\partial_\tau S(\mathbf{r}, t) = -S + d\partial^2 S + \sum_i h(\mathbf{X}_i)\delta(\mathbf{r} - \mathbf{r}_i), \quad (7)$$

where  $\mathbf{X}_i$  represents the internal state (which will later be identified as a phase),  $\mathbf{r}$  represents the position of the  $i$ -th oscillator, and  $S$  represents the concentration of the background chemical. By means of a center manifold calculation and a phase reduction technique they derived the simpler equations

$$\dot{\psi}_i(t) = \sum_{j \neq i} e^{-|\mathbf{R}_{ji}|} \sin(\Psi_{ji} - \alpha|\mathbf{R}_{ji}| - c_1) \quad (8)$$

$$\dot{\mathbf{r}}_i(t) = c_3 \sum_{j \neq i} \hat{\mathbf{R}}_{ji} e^{-|\mathbf{R}_{ji}|} \sin(\Psi_{ji} - \alpha|\mathbf{R}_{ji}| - c_2) \quad (9)$$

where  $\mathbf{R}_{ji} = \mathbf{R}_j - \mathbf{R}_i$ ,  $\Psi_{ji} = \psi_j - \psi_i$ , and  $\psi_i$  is the  $i$ -th oscillator's phase. We call this Iwasa-Tanaka model. Notice the space-phase coupling in this model is somewhat peculiar; in contrast to the swarmalator model given by (3) and (4) the relative position  $\mathbf{R}_{ji}$  and relative phase  $\Psi_{ji}$  appear *inside* the sine terms in both the  $\dot{\mathbf{r}}_i$  and  $\dot{\psi}_i$  equations. Another difference between the two models is that  $\dot{\mathbf{r}}_i$  in (8) has no hardshell repulsion term, which means the oscillators can occupy the same position in space.

The Iwasa-Tanaka model has rich collective states. An exhaustive catalogue of these states with respect to the model's four parameters is an ongoing effort.<sup>57</sup> Highlights include a family of clustered states<sup>44,45,56</sup> in which swarmalators collect in synchronous groups. The spatial distributions of these groups depend on their phase, similar to the splintered phase wave (Figure 1). The authors speculate this phase clustering is reminiscent of the chemotactic cell sorting during biological development.<sup>44</sup> The Iwasa-Tanaka model also produces ring states,<sup>45</sup> as well as an interesting 'juggling' state<sup>54</sup> in which the population forms a "rotating triangular structure whose corners appear to 'catch' and 'throw' individual elements" – in other words, the population juggles the elements around the corners of a triangle (see Figure 1 in<sup>54</sup>). Aside from theoretical novelty, this juggling could conceivably be exploited in robotic swarms, potentially allowing some form of relay between the elements.

## 2.2 Alignment and synchronization

In our proposed taxonomy, systems that combine alignment and synchronization are characterized by an internal phase  $\theta$  and an orientation  $\beta$  without a dynamic spatial degree of freedom. In other words, the particles' position  $\mathbf{x}$  might affect their  $\theta$  and  $\beta$  dynamics, but  $\mathbf{x}$  itself does not evolve in time. Although units characterized by just a phase  $\theta$  and orientation  $\beta$  might seem odd, Leon and Liverpool studied a system with units which meet these criteria: a class of soft active fluids which constitute a 'new type of nonequilibrium soft matter – a space-time liquid crystal'.<sup>58</sup> They developed a phenomenological theory of these soft fluids which allowed them to derive dynamical equations for order parameters quantifying the orientation order, phase order, and orientation-phase order. These revealed collective states which maximize each of these order parameters: aligned states with orientational order but no phase order, sync'd states with phase order but no orientational order, and states with both phase and orientational order. They were able to partially analyze these states, and conjectured the states could be realized in protein-filaments mixtures, such as cell cytoskeletons<sup>59,60</sup> or tissue-forming cells.<sup>61</sup>

## 2.3 Alignment, aggregation, and synchronization

Systems with units that align, aggregate, and synchronize – and therefore have dynamic state variables  $\mathbf{x}, \theta, \beta$  – are the least well studied. A small study was carried out in,<sup>48</sup> with the aim of checking if the swarmalator states reported are generic, i.e., robust to the inclusion of alignment dynamics (it was found they were). Beyond this preliminary study, the space of possible behaviors arising from alignment, aggregation, and synchronization is largely unexplored.

## 2.4 Alignment and aggregation

While not strictly within our proposed taxonomy of swarmalator systems, we give a brief review of studies on aggregation and alignment. We do this because alignment can be viewed as a type of synchronization, where instead of units adjusting to a common phase  $\theta$ , units instead adjust a common orientation  $\beta$ . Or put another way, because an internal phase  $\theta$  and orientation  $\beta$  are formally equivalent – both being circular variables – particles aligning their orientations is analogous to oscillators aligning their phases.

Vicsek set the paradigm of this class of models with a beautiful and simple model:

$$\mathbf{r}_i(t + \delta t) = \mathbf{r}_i(t) + (\nu \delta t) \hat{n} \quad (10)$$

$$\beta_i(t + \delta t) = \langle \beta_j \rangle_{|\mathbf{r}_j - \mathbf{r}_i| < r} + \eta_i(t), \quad (11)$$

where  $\hat{n} = (\cos(\beta), \sin(\beta))$ ,  $\eta_i(t)$  is a white noise,  $\nu$  is the particles' speed, and  $r$  is the coupling range. For sufficiently strong noise this model exhibits a flocking transition, where particles switch from erratic incoherent movements to moving in a unidirectional flock.

Leibchen and Levis<sup>62,63</sup> modified the Vicsek model as follows:

$$\dot{\mathbf{r}}_i = \nu \vec{n}_i \tag{12}$$

$$\dot{\beta}_i = \omega_i + \frac{K}{\pi R^2} \sum_{j \in \partial_i} \sin(\theta_j - \theta_i) + \sqrt{2D} \eta_i, \tag{13}$$

where  $\partial_i$  is the set of neighbours of the  $i$ -th particle,  $K$  is the phase coupling, and  $D$  is the noise strength. The key new feature is the  $\omega_i$  term, signifying the particles have an intrinsic rotation. They find three collective states: a disordered state in which neither spatial order nor phase order exists, a chiral state in which macroscopic 'droplets' of synchronized oscillators form in a sea of otherwise desynchronized oscillators, and finally a *mutual flocking* state in which oscillators with 'opposite chirality cooperate to move coherently at a relative angle, forming non-rotating flocks'. The theoretical novelty of the latter state is that – in contrast to regular, non-moving, locally-coupled oscillators coupled in two or three dimensions – long-range synchronous clusters are observed, despite the fact that the oscillators are only locally coupled.

### 3. SWARMALATORS IN THE REAL-WORLD

To our knowledge, there are just two works which unequivocally realize swarmalators in the real world – by this we mean, precisely, a real-world system in which a bidirectional space-phase coupling is unambiguously exhibited. These works are:

**Swarmalatorbots.** Bettstetter et al first realized the collective states of the swarmalator model (Figure 1) in the lab using small robots.<sup>64</sup> They programmed 'swarmalatorbots' whose governing equations were derived from the swarmalator model (3), (4).

**Magnetic domain walls.** To our knowledge, the first realization of a natural swarmalator system was found by Hrabec et al when studying the magnetic domain walls.<sup>65</sup> Ordinarily, domain walls are described by a single spatial degree of freedom  $\mathbf{x}$ . But as the authors note, the dynamics on the walls also depend on their internal structure. The authors minimally describe this internal state by a one dimensional polarization angle of the internal magnetic field  $\theta$ . This allows the wall to be viewed as a point particle with position  $\mathbf{x}$  and phase  $\theta$  – to be viewed as a swarmalator.<sup>65</sup> An experimental study of coupling between two domain walls – usually just one wall is studied – revealed the walls can synchronize, which in turn affects the walls' velocity. Richer space-phase dynamics, such as families of Lissajous curves, are also reported.

\*\*\*

Beyond swarmalatorbots and magnetic domain walls, there are many systems in which a bidirectional coupling between swarming and synchronization might exist. We list these candidate swarmalator systems below:

**Myxobacteria.** Myxobacteria are bacteria commonly found in soil, and can produce interesting collective effects.<sup>66</sup> The phase variable of myxobacteria characterizes the internal growth cycle of the cell. Groups of cells interact through cell-to-cell contact, theorized to provide a channel through which swarming and synchronization can couple. Populations of myxobacteria can exhibit a 'ripple phase', in which complex patterns of waves propagate through population.<sup>66</sup> In contrast to other wave phenomena in biological systems, such as the well-studied *Dictyostelium discoideum*, these rippling waves do not annihilate on collision. In<sup>66</sup> a Fokker-Planck type equation was used to analyze rippling waves. Realizing them in a microscopic swarmalator system is an open problem.

**Biological microswimmers.** 'Microswimmers' is an umbrella term for self-propelled microorganisms confined to fluids, such as cilia, bacteria, and sperm.<sup>67</sup> Groups of microswimmers show swarming behavior as a result of cooperative goal seeking, such as searching for food or light. They can also synchronize: Here the



phase variable is associated with the rhythmic beating of swimmers' tails which – through hydrodynamic interactions – can synchronize with the beatings of others swimmers' tails. Whether or not this hydrodynamics provides a bidirectional coupling between sync and swarming – as required of swarmalators – is unclear, but to us seems plausible. Researchers have developed models of sperm under this assumption<sup>68</sup> and found clusters of synchronized sperm consistent with real data.<sup>69</sup> Vortex arrays of sperm have also been reported.<sup>70</sup> Here, sperm self-organize into subgroups arranged in a lattice; within each subgroup, the sperm move in a vortex, wherein a correlation between their angular velocity and their phase velocity is realized, reminiscent of the splintered phase wave state (Figure 1). Simulations of realistic models have been carried out<sup>47</sup> which show other interesting phenomena.

**Japanese tree frogs.** During mating season, male Japanese tree frogs attract females by croaking rhythmically. The croaking of neighbouring frogs tend to anti-synchronize due to a precedence effect: croaking shortly after a rival makes a frog look less dominant. Researchers have theorized that this competition leads to mutual interaction between frogs' space and phase dynamics, coupling sync and swarming. Models based on these assumption produce ring-like states where frogs arrange themselves on the borders of fields with interesting space-phase patterns, some of which are consistent with data on real tree-frogs' behavior collected in the wild.

**Magnetic colloids.** Similar to magnetic domain walls, a colloid's constituent particles can synchronize the orientation of their magnetic dipole vector when sufficiently close. When in solution, they are free to move around in space, creating a feedback loop between their space and phase dynamics. In ferromagnetic colloids confined to liquid-liquid interfaces, Snezhko and Aranson<sup>71</sup> found this interaction leads to the formation of 'asters' – star-like arrangements of particles whose spatial angles correlate with the orientations of their magnetic dipole vectors, equivalent to the static phase wave (Figure 1). Yan et al explored how synchronization is useful in colloids of Janus particles.<sup>72</sup> Janus particles are micrometer-sized spheres with one hemisphere covered with nickle which gives them non-standard magnetic properties. In particular when subject to a precessing magnetic field, they oscillate about their centers of mass. This oscillation creates a coupling between particles, giving rise to 'synchronization-selected' self-assembly. For example, zig-zag chains of particles and microtubes of synchronized particles can be realized.

#### 4. SWARMALATORS IN BIO-INSPIRED COMPUTING

The computer engineering community has been inspired by coupled oscillators to develop new techniques for synchronization in communication and sensor networks.<sup>21–24,73,74</sup> Here, instead of Kuramoto-type oscillators (2), models of pulse-coupled oscillators have been borrowed. As the name suggests, pulse-coupled oscillators communicate by exchanging short signals; this time-discrete variant of coupling is a more natural fit for application in technology, where smooth, continuous coupling – as exemplified by Kuramoto oscillators – is costly to achieve. The canonical model of pulse-coupled oscillators is the Peskin model,<sup>17</sup> defined by

$$\dot{x}_i = S_0 - \gamma x_i \tag{14}$$

with  $S_0, \gamma > 0$  and  $x_i$  is a voltage like state variable for the  $i$ -th oscillator. When the oscillators' voltage reach a threshold value they ( $i$ ) fire a pulse which instantaneously raises the voltage of all the other oscillators ( $ii$ ) reset their voltage to zero, along with any other oscillator whose voltage was raised above the threshold on account of receiving a pulse. The synchronization properties of the Peskin model are well-studied.<sup>75,76</sup> More recently, estimates for the convergence speed have been derived.<sup>77,78</sup>

The simplicity, distributed nature, adaptability, and scalability of pulse-coupled oscillators make them attractive from an engineering point of view, where temporal coordination is often a goal. For example, synchronization is required for many tasks in different layers of computing and communications systems<sup>79</sup> such as in the alignment of transmission slots for efficient medium access,<sup>80</sup> scheduling of sleep cycles for energy efficiency,<sup>81</sup> and coordination of sensor readings to capture a scene from different perspectives.<sup>82</sup> It is the relative synchrony of the networked units, not necessarily the absolute time, that is relevant here. The synchronization precision required is determined by the specific task. The achievable precision depends on the environment and hardware, and is influenced by deviations in delays and phase rates. Experiments in laboratory environments show that the precision is in the order of hundred microseconds with low-cost programmable sensor platforms<sup>21,83</sup> and a few



microseconds with field-programmable gate array (FPGA)-based radio boards.<sup>84</sup> Further use cases can be found in acoustics (synchronizing multiple loudspeakers) and energy systems (synchronizing decentralized grids<sup>85,86</sup>), to give two examples.

Synchronization is also important in robotics in order to perform coordinated movements – and this is where temporal and spatial coordination unite. As mentioned, Bettstetter and colleagues<sup>64</sup> extended the swarmalator model<sup>48</sup> so that it could be applied to mobile robots. They implemented the extended model in the Robot Operating System 2 (ROS 2) and experimentally demonstrated that the space-time patterns achieved in theory (Figure 1) can be reproduced in the real world. Beyond realizing these specific states, we conjecture that swarmalator-type models will enable novel self-assembly procedures in other robotics groups, in turn enabling collaborative actions in monitoring, exploration, and manipulation. As envisioned in,<sup>64</sup> underwater robots could be designed, which – in imitation of biological microswimmers – could both swim in formation and synchronize their fin movements, thereby enhancing their functional capabilities.

More speculatively, the swarmalator concept could be useful in the design of autonomous transport systems; for example, when multiple vehicles driving in a convoy have to avoid collisions with other convoys due to crossings and for the purpose of overtaking. The model could also enable self-configuring distributed antenna arrays (or loudspeakers) in which multiple antenna elements (or sound sources) automatically arrange their positions and orientations to create specific radiation patterns used to send radio (audio) signals in a synchronous way. Another promising application field is the planning and replanning of processes in factories, where products and machines must follow a certain space-time order and where standard optimization methods reach their limits due to the high system complexity. A final, more playful, application is in art: Artistic aerial light shows created by drone swarms running the swarmalator model produce charming visual displays.

## 5. DISCUSSION

The implicit strategy in the theoretical works we reviewed is: given a model, which spatiotemporal patterns emerge? Future work could consider the inverse strategy: given a desired spatiotemporal pattern, which form should the model take on? – a question often considered in engineering contexts. From a more general perspective, one has to design the local rules and interactions that guide a self-organizing system toward a desired global state.<sup>87</sup> Von Brecht et al have studied this inverse question for swarming systems;<sup>88</sup> perhaps their tools could be extended to swarmalators.

Future work could also extend the reviewed models, which – recall – were designed to be as minimal as possible. A wealth of synchronization phenomena have been found by adorning the Kuramoto model with new features like mixed-sign coupling,<sup>89–92</sup> non-local coupling,<sup>11,12</sup> and delayed interactions.<sup>93</sup> Equipping swarmalators with these features would likely cure the poverty of phenomena in the first, third, and fourth quadrants of the  $(J, K)$  plane (Figure 2) and produce new dynamics in Iswasa-tanaka and Vicsek type models, too. Phase models more sophisticated than the Kuramoto model could also be explored; the Winfree model<sup>94</sup> or the newly introduced Janus oscillators<sup>95</sup> would be exciting to experiment with. Swarmalators with discrete and spatially local coupling also merit study and are useful in computer and communication systems. Stitching the Peskin model (14) to the aggregation equation (1) or the Vicsek model (11) seems the natural way to do this. Continuing to study finite systems of swarmalators<sup>53</sup> is also pertinent, since many robotic systems lie in the low  $N$  regime. And finally, beyond richer phase models, the spatial dynamics of swarmalators could also be generalized. Even within the framework of the aggregation equation (1), a menagerie of spatial patterns have been catalogued.<sup>96</sup> Swarmalator counterparts to these states would be exciting to explore.

We hope to have outlined that swarmalators have great potential in computing and other fields of technology. Here we see swarmalators only as a case study in a broader class of systems with large technological utility: systems whose units have both spatial and internal degrees of freedom. The one-dimensional phase  $\theta$  of a swarmalator is perhaps the simplest instance of an internal degree of freedom, which can more generally be represented by a feature vector  $\mathbf{f}$ . This vector could describe a particle's (three dimensional) magnetic or electric field; a person's political affiliation, mood, or health state; a bacteria's phase in a non-circular developmental cycle; and the mode of a robot, machine, or vehicle. The collective states arising when particles' features  $\mathbf{f}$  and positions  $\mathbf{x}$  defines a wide, scarcely investigated area. We anticipate an exploration of this area will bear rich fruit in both Nature and technology.

## Acknowledgements

The work of C. Bettstetter is supported by the Austrian Science Fund (FWF) [grant P30012] and the Popper Kolleg *Networked Autonomous Aerial Vehicles*.

The work of Kevin O’Keeffe thanks Allianz, Amsterdam Institute for Advanced Metropolitan Solutions, Brose, Cisco, Ericsson, Fraunhofer Institute, Liberty Mutual Institute, Kuwait-MIT Center for Natural Resources and the Environment, Shenzhen, Singapore- MIT Alliance for Research and Technology (SMART), UBER, Vitoria State Government, Volkswagen Group America, and all the members of the MIT Senseable City Lab Consortium for supporting this research.

## REFERENCES

- [1] Winfree, A. T., [*The geometry of biological time*], vol. 12, Springer Science & Business Media (2001).
- [2] Kuramoto, Y., “International symposium on mathematical problems in theoretical physics,” *Lecture Notes in Physics* **30**, 420 (1975).
- [3] Strogatz, S., [*Sync: The emerging science of spontaneous order*], Penguin UK (2004).
- [4] Pikovsky, A., Rosenblum, M., and Kurths, J., [*Synchronization: a universal concept in nonlinear sciences*], vol. 12, Cambridge University Press (2003).
- [5] Acebrón, J. A., Bonilla, L. L., Vicente, C. J. P., Ritort, F., and Spigler, R., “The kuramoto model: A simple paradigm for synchronization phenomena,” *Reviews of Modern Physics* **77**(1), 137 (2005).
- [6] Daido, H., “Quasientrainment and slow relaxation in a population of oscillators with random and frustrated interactions,” *Physical Review Letters* **68**(7), 1073 (1992).
- [7] Bonilla, L., Vicente, C. P., and Rubi, J., “Glassy synchronization in a population of coupled oscillators,” *Journal of Statistical Physics* **70**(3-4), 921–937 (1993).
- [8] Iatsenko, D., McClintock, P. V., and Stefanovska, A., “Glassy states and super-relaxation in populations of coupled phase oscillators,” *Nature Communications* **5**, 4118 (2014).
- [9] Ottino-Löffler, B. and Strogatz, S. H., “Volcano transition in a solvable model of frustrated oscillators,” *Physical Review Letters* **120**(26), 264102 (2018).
- [10] Kuramoto, Y. and Battogtokh, D., “Coexistence of coherence and incoherence in nonlocally coupled phase oscillators,” *arXiv preprint cond-mat/0210694* (2002).
- [11] Abrams, D. M. and Strogatz, S. H., “Chimera states for coupled oscillators,” *Physical Review Letters* **93**(17), 174102 (2004).
- [12] Abrams, D. M., Mirollo, R., Strogatz, S. H., and Wiley, D. A., “Solvable model for chimera states of coupled oscillators,” *Physical Review Letters* **101**(8), 084103 (2008).
- [13] Montbrió, E., Pazó, D., and Roxin, A., “Macroscopic description for networks of spiking neurons,” *Physical Review X* **5**(2), 021028 (2015).
- [14] Pazó, D. and Montbrió, E., “Low-dimensional dynamics of populations of pulse-coupled oscillators,” *Physical Review X* **4**(1), 011009 (2014).
- [15] O’Keeffe, K. P. and Strogatz, S. H., “Dynamics of a population of oscillatory and excitable elements,” *Physical Review E* **93**(6), 062203 (2016).
- [16] Luke, T. B., Barreto, E., and So, P., “Complete classification of the macroscopic behavior of a heterogeneous network of theta neurons,” *Neural Computation* **25**(12), 3207–3234 (2013).
- [17] Peskin, C. S., “Mathematical aspects of heart physiology,” *Courant Inst. Math* (1975).
- [18] Grudziński, K. and Żebrowski, J. J., “Modeling cardiac pacemakers with relaxation oscillators,” *Physica A: Statistical Mechanics and its Applications* **336**(1-2), 153–162 (2004).
- [19] Gois, S. R. and Savi, M. A., “An analysis of heart rhythm dynamics using a three-coupled oscillator model,” *Chaos, Solitons & Fractals* **41**(5), 2553–2565 (2009).
- [20] Saw, V.-L., Chung, N. N., Quek, W. L., Pang, Y. E. I., and Chew, L. Y., “Bus bunching as a synchronisation phenomenon,” *arXiv preprint arXiv:1812.00609* (2018).

- [21] Werner-Allen, G., Tewari, G., Patel, A., Welsh, M., and Nagpal, R., “Firefly-inspired sensor network synchronicity with realistic radio effects,” in [*Proc. ACM Conf. Embedded Networked Sensor Systems (SenSys)*], (Nov. 2005).
- [22] Hong, Y.-W. and Scaglione, A., “A scalable synchronization protocol for large scale sensor networks and its applications,” *IEEE J. Select. Areas Commun.* **23**(5), 1085–1099 (2005).
- [23] Babaoglu, O., Binci, T., Jelasity, M., and Montresor, A., “Firefly-inspired heartbeat synchronization in overlay networks,” in [*Proc. Intern. Conf. on Self-Adaptive and Self-Organizing Systems (SASO)*], 77–86, IEEE (2007).
- [24] Tyrrell, A., Auer, G., and Bettstetter, C., “Emergent slot synchronization in wireless networks,” *IEEE Trans. Mobile Comput.* **9**, 719–732 (May 2010).
- [25] Vicsek, T., Czirók, A., Ben-Jacob, E., Cohen, I., and Shochet, O., “Novel type of phase transition in a system of self-driven particles,” *Physical Review Letters* **75**(6), 1226 (1995).
- [26] Ramaswamy, S., “The mechanics and statistics of active matter,” *Annu. Rev. Condens. Matter Phys.* **1**(1), 323–345 (2010).
- [27] Marchetti, M. C., Joanny, J.-F., Ramaswamy, S., Liverpool, T. B., Prost, J., Rao, M., and Simha, R. A., “Hydrodynamics of soft active matter,” *Reviews of Modern Physics* **85**(3), 1143 (2013).
- [28] Sanchez, T., Chen, D. T., DeCamp, S. J., Heymann, M., and Dogic, Z., “Spontaneous motion in hierarchically assembled active matter,” *Nature* **491**(7424), 431 (2012).
- [29] Dorigo, M. and Birattari, M., [*Ant colony optimization*], Springer (2010).
- [30] Meng, X., Liu, Y., Gao, X., and Zhang, H., “A new bio-inspired algorithm: chicken swarm optimization,” in [*Proc. Intern. Conf. in Swarm Intelligence*], 86–94, Springer (2014).
- [31] Meng, X.-B., Gao, X. Z., Lu, L., Liu, Y., and Zhang, H., “A new bio-inspired optimisation algorithm: Bird swarm algorithm,” *Journal of Experimental & Theoretical Artificial Intelligence* **28**(4), 673–687 (2016).
- [32] Mirjalili, S., Gandomi, A. H., Mirjalili, S. Z., Saremi, S., Faris, H., and Mirjalili, S. M., “Salp swarm algorithm: A bio-inspired optimizer for engineering design problems,” *Advances in Engineering Software* **114**, 163–191 (2017).
- [33] Yang, X.-S., Cui, Z., Xiao, R., Gandomi, A. H., and Karamanoglu, M., [*Swarm intelligence and bio-inspired computation: theory and applications*], Newnes (2013).
- [34] Shi, Y. et al., “Particle swarm optimization: developments, applications and resources,” in [*Proc. Congress on Evolutionary Computation*], **1**, 81–86, IEEE (2001).
- [35] Hamann, H., [*Swarm Robotics: A Formal Approach*], Springer (2018).
- [36] Schmickl, T. and Hamann, H., “Beeclust: A swarm algorithm derived from honeybees,” *Bio-inspired computing and communication networks*, 95–137 (2011).
- [37] Savkin, A. V., “Coordinated collective motion of groups of autonomous mobile robots: Analysis of vicsek’s model,” *IEEE Transactions on Automatic Control* **49**(6), 981–982 (2004).
- [38] Merkle, D., Middendorf, M., and Scheidler, A., “Swarm controlled emergence-designing an anti-clustering ant system,” in [*Proc. IEEE Swarm Intelligence Symp.*], 242–249 (2007).
- [39] von Brecht, J. H. and Uminsky, D. T., “Anisotropic assembly and pattern formation,” *Nonlinearity* **30**(1), 225 (2016).
- [40] Uriu, K., Ares, S., Oates, A. C., and Morelli, L. G., “Dynamics of mobile coupled phase oscillators,” *Physical Review E* **87**(3), 032911 (2013).
- [41] Sevilla, F. J., Dossetti, V., and Heiblum-Robles, A., “Synchronization and collective motion of globally coupled brownian particles,” *Journal of Statistical Mechanics: Theory and Experiment* **2014**(12), P12025 (2014).
- [42] Stilwell, D. J., Boltt, E. M., and Roberson, D. G., “Sufficient conditions for fast switching synchronization in time-varying network topologies,” *SIAM Journal on Applied Dynamical Systems* **5**(1), 140–156 (2006).
- [43] Frasca, M., Buscarino, A., Rizzo, A., Fortuna, L., and Boccaletti, S., “Synchronization of moving chaotic agents,” *Physical Review Letters* **100**(4), 044102 (2008).
- [44] Tanaka, D., “General chemotactic model of oscillators,” *Physical Review Letters* **99**(13), 134103 (2007).
- [45] Iwasa, M., Iida, K., and Tanaka, D., “Hierarchical cluster structures in a one-dimensional swarm oscillator model,” *Physical Review E* **81**(4), 046220 (2010).

- [46] Starnini, M., Frasca, M., and Baronchelli, A., “Emergence of metapopulations and echo chambers in mobile agents,” *Scientific reports* **6**, 31834 (2016).
- [47] Belovs, M., Livanovičs, R., and Cēbers, A., “Synchronized rotation in swarms of magnetotactic bacteria,” *Physical Review E* **96**(4), 042408 (2017).
- [48] O’Keeffe, K. P., Hong, H., and Strogatz, S. H., “Oscillators that sync and swarm,” *Nature Communications* **8**(1), 1504 (2017).
- [49] Topaz, C. M. and Bertozzi, A. L., “Swarming patterns in a two-dimensional kinematic model for biological groups,” *SIAM Journal on Applied Mathematics* **65**(1), 152–174 (2004).
- [50] Mogilner, A. and Edelstein-Keshet, L., “A non-local model for a swarm,” *Journal of Mathematical Biology* **38**(6), 534–570 (1999).
- [51] Kuramoto, Y., “Collective synchronization of pulse-coupled oscillators and excitable units,” *Physica B* **50**, 15–30 (1991).
- [52] Hong, H., “Active phase wave in the system of swarms with attractive phase coupling,” *Chaos: An Interdisciplinary Journal of Nonlinear Science* **28**(10), 103112 (2018).
- [53] O’Keeffe, K. P., Evers, J. H., and Kolokolnikov, T., “Ring states in swarms with attractive coupling,” *Physical Review E* **98**(2), 022203 (2018).
- [54] Iwasa, M., Iida, K., and Tanaka, D., “Juggling motion in a system of motile coupled oscillators,” *Physical Review E* **83**(3), 036210 (2011).
- [55] Iwasa, M. and Tanaka, D., “Mechanism underlying the diverse collective behavior in the swarm oscillator model,” *Physics Letters A* **381**(36), 3054–3061 (2017).
- [56] Iwasa, M. and Tanaka, D., “Dimensionality of clusters in a swarm oscillator model,” *Physical Review E* **81**(6), 066214 (2010).
- [57] Iwasa, M., Iida, K., and Tanaka, D., “Various collective behavior in swarm oscillator model,” *Physics Letters A* **376**(30-31), 2117–2121 (2012).
- [58] Leoni, M. and Liverpool, T., “Synchronization and liquid crystalline order in soft active fluids,” *Physical Review Letters* **112**(14), 148104 (2014).
- [59] Toner, J. and Tu, Y., “Long-range order in a two-dimensional dynamical xy model: how birds fly together,” *Physical Review Letters* **75**(23), 4326 (1995).
- [60] Ahmadi, A., Marchetti, M. C., and Liverpool, T. B., “Hydrodynamics of isotropic and liquid crystalline active polymer solutions,” *Physical Review E* **74**(6), 061913 (2006).
- [61] Sachs, T., [*Pattern formation in plant tissues*], vol. 25, Cambridge University Press (2005).
- [62] Liebchen, B. and Levis, D., “Collective behavior of chiral active matter: pattern formation and enhanced flocking,” *Physical Review Letters* **119**(5), 058002 (2017).
- [63] Levis, D., Pagonabarraga, I., and Liebchen, B., “Activity induced synchronization: From mutual flocking to chiral self-sorting,” *arXiv preprint arXiv:1802.02371* (2018).
- [64] Gniewek, A., Barciś, M., and Bettstetter, C., “Robots that sync and swarm: A proof of concept in ROS 2,” *arXiv preprint arXiv:1903.06440* (2019).
- [65] Hrabec, A., Krížáková, V., Pizzini, S., Sampaio, J., Thiaville, A., Rohart, S., and Vogel, J., “Velocity enhancement by synchronization of magnetic domain walls,” *Physical Review Letters* **120**(22), 227204 (2018).
- [66] Igoshin, O. A., Mogilner, A., Welch, R. D., Kaiser, D., and Oster, G., “Pattern formation and traveling waves in myxobacteria: theory and modeling,” *Proceedings of the National Academy of Sciences* **98**(26), 14913–14918 (2001).
- [67] Elgeti, J., Winkler, R. G., and Gompper, G., “Physics of microswimmers: single particle motion and collective behavior: a review,” *Reports on progress in physics* **78**(5), 056601 (2015).
- [68] Yang, Y., Elgeti, J., and Gompper, G., “Cooperation of sperm in two dimensions: synchronization, attraction, and aggregation through hydrodynamic interactions,” *Physical Review E* **78**(6), 061903 (2008).
- [69] Hayashi, F., “Insemination through an externally attached spermatophore: bundled sperm and post-copulatory mate guarding by male fishflies (megalopectera: Corydalidae),” *Journal of Insect Physiology* **42**(9), 859–866 (1996).
- [70] Riedel, I. H., Kruse, K., and Howard, J., “A self-organized vortex array of hydrodynamically entrained sperm cells,” *Science* **309**(5732), 300–303 (2005).

- [71] Snezhko, A. and Aranson, I. S., “Magnetic manipulation of self-assembled colloidal asters,” *Nature Materials* **10**(9), 698 (2011).
- [72] Yan, J., Bloom, M., Bae, S. C., Luijten, E., and Granick, S., “Linking synchronization to self-assembly using magnetic janus colloids,” *Nature* **491**(7425), 578 (2012).
- [73] Mathar, R. and Mattfeldt, J., “Pulse-coupled decentral synchronization,” *SIAM J. Appl. Math.* **56**(4), 1094–1106 (1996).
- [74] Simeone, O., Spagnolini, U., Bar-Ness, Y., and Strogatz, S. H., “Distributed synchronization in wireless networks,” *IEEE Signal Processing Mag.* **25**, 81–97 (Sept. 2008).
- [75] Bottani, S., “Pulse-coupled relaxation oscillators: from biological synchronization to self-organized criticality,” *Physical Review Letters* **74**(21), 4189 (1995).
- [76] Mirollo, R. E. and Strogatz, S. H., “Synchronization of pulse-coupled biological oscillators,” *SIAM Journal on Applied Mathematics* **50**(6), 1645–1662 (1990).
- [77] O’Keeffe, K. P., Krapivsky, P. L., and Strogatz, S. H., “Synchronization as aggregation: Cluster kinetics of pulse-coupled oscillators,” *Physical Review Letters* **115**(6), 064101 (2015).
- [78] O’Keeffe, K. P., “Transient dynamics of pulse-coupled oscillators with nonlinear charging curves,” *Physical Review E* **93**(3), 032203 (2016).
- [79] Bregni, S., [*Synchronization of Digital Telecommunications Networks*], Wiley, New York, NY, USA (2002).
- [80] Roberts, L. G., “Aloha packet system with and without slots and capture,” *SIGCOMM Comput. Commun. Rev.* **5**, 28–42 (Apr. 1975).
- [81] Ye, W., Heidemann, J., and Estrin, D., “Medium access control with coordinated adaptive sleeping for wireless sensor networks,” *IEEE/ACM Trans. Netw.* **12**, 493–506 (June 2004).
- [82] Aghajan, H. and Cavallaro, A., [*Multi-Camera Networks: Principles and Applications*], Academic Press (2009).
- [83] Pagliari, R. and Scaglione, A., “Scalable network synchronization with pulse-coupled oscillators,” *IEEE Trans. Mobile Comput.* **10**(3), 392–405 (2011).
- [84] Brandner, G., Schilcher, U., and Bettstetter, C., “Firefly synchronization with phase rate correction and its experimental analysis in wireless systems,” *Computer Networks* **97**, 74–87 (Mar. 2016).
- [85] Rohden, M., Sorge, A., Timme, M., and Witthaut, D., “Self-organized synchronization in decentralized power grids,” *Physical Review Letters* **109**(6), 064101 (2012).
- [86] Skardal, P. S. and Arenas, A., “Control of coupled oscillator networks with application to microgrid technologies,” *Science Advances* **1**(7), e1500339 (2015).
- [87] Prehofer, C. and Bettstetter, C., “Self-organization in communication networks: Principles and design paradigms,” *IEEE Commun. Mag.* **43**, 78–85 (July 2005).
- [88] von Brecht, J. H. and Uminsky, D., “On soccer balls and linearized inverse statistical mechanics,” *Journal of Nonlinear Science* **22**(6), 935–959 (2012).
- [89] Hong, H. and Strogatz, S. H., “Kuramoto model of coupled oscillators with positive and negative coupling parameters: an example of conformist and contrarian oscillators,” *Physical Review Letters* **106**(5), 054102 (2011).
- [90] Maistrenko, Y., Penkovsky, B., and Rosenblum, M., “Solitary state at the edge of synchrony in ensembles with attractive and repulsive interactions,” *Physical Review E* **89**(6), 060901 (2014).
- [91] Hong, H., O’Keeffe, K. P., and Strogatz, S. H., “Phase coherence induced by correlated disorder,” *Physical Review E* **93**(2), 022219 (2016).
- [92] Hong, H., O’Keeffe, K. P., and Strogatz, S. H., “Correlated disorder in the kuramoto model: Effects on phase coherence, finite-size scaling, and dynamic fluctuations,” *Chaos: An Interdisciplinary Journal of Nonlinear Science* **26**(10), 103105 (2016).
- [93] Yeung, M. S. and Strogatz, S. H., “Time delay in the kuramoto model of coupled oscillators,” *Physical Review Letters* **82**(3), 648 (1999).
- [94] Winfree, A. T., “Biological rhythms and the behavior of populations of coupled oscillators,” *J. Theoretical Biology* **16**, 15–42 (July 1967).
- [95] Nicolaou, Z. G., Eroglu, D., and Motter, A. E., “Multifaceted dynamics of janus oscillator networks,” *Physical Review X* **9**(1), 011017 (2019).
- [96] Kolokolnikov, T., Sun, H., Uminsky, D., and Bertozzi, A. L., “Stability of ring patterns arising from two-dimensional particle interactions,” *Physical Review E* **84**(1), 015203 (2011).